

COVID Exposure Assessment Tool (CEAT) - US Units

A Simple-to-Use Tool (BETA Version) for Group Exposure to SARS-COV-2 based upon Mechanistic Factors, Group Behavior, and Infection Prevalence in the Community. **Open with Adobe Reader® or Adobe Acrobat®.**

Start Step 1 Select the factor from the table below that best matches the group in the activity.

| The group is composed of people who... | Group Factor |
|--|--------------|
| Are known to be following all public health guidance. | 1 |
| You think are following all public health guidance. | 17 |
| Who are from the general public. | 50 |
| Includes healthcare workers in contact with diagnosed COVID-19 patients. | 58 |
| You suspect are not compliant with public health guidance. | 67 |
| Are known to be diagnosed with active COVID-19 | 100 |


Scenario A Scenario B

Group Factor Adjustment: A: B:

Step 2 Enter the number of people sharing the space for the activity. Must be between 2 and 250 people.

Number of People Sharing Activity Space: A: B:

Step 3 Select Distancing Adjustment



Distancing Adjustment: A: B:

Step 4 Select Mask Effectiveness Adjustment

| | 4-1 Exhalation % of People Wearing Masks | | | | 4-2 Inhalation % of People Wearing Masks | | | |
|---------------|---|-----|-----|-----|---|-----|-----|-----|
| | ALL | 75% | 50% | 25% | ALL | 90% | 50% | 25% |
| N95/KN95 | -22 | -14 | -5 | -2 | -17 | -12 | -4 | -2 |
| Surgical Mask | -12 | -9 | -4 | -2 | -8 | -6 | -3 | -1 |
| Average Mask | -5 | -4 | -2 | -1 | -5 | -4 | -2 | -1 |
| Cloth Mask | -5 | -4 | -2 | -1 | -3 | -2 | -1 | -1 |
| No Mask | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

Mask Effectiveness Adjustment: A: B: A: B:

Step 5 Select Vocalization Intensity Adjustment

| Activity | Exhalation Type | Adjust |
|----------------|-----------------|--------|
| Resting | Silent/ Sleep | -13 |
| | Speaking | -1 |
| Standing | Loudly speaking | +12 |
| | Silent | -12 |
| | Speaking | 0 |
| Light Exertion | Singing | +18 |
| | Silent | -5 |
| | Speaking | +6 |
| Heavy Exertion | Loudly speaking | +20 |
| | Silent | +1 |
| Heavy Exertion | Speaking | +12 |
| | Loudly speaking | +26 |

Vocalization Adjustment:

A: B:

Step 6 Select Breathing Rate Adjustment

| Activity | Adjust |
|----------------|--------|
| Sleep | -8 |
| Resting | -7 |
| Passive | 0 |
| Light Exertion | +6 |
| Heavy Exertion | +10 |

Breathing Adjustment:

A: B:

Step 7 Enter the duration that most closely matches activity.

Duration of Activity in Hours: A: B:

Step 8 Select whether outdoor or indoor and the matching Ventilation Adjustment
A: B: Outdoor activities: Select wind conditions that best match.

Outdoor?

| Beaufort Scale | | | Adjust |
|----------------|-------------|--|--------|
| Very Calm | 0.1 - 1 mph | No direction or flow observed | -16 |
| Calm | 1 - 3 mph | Direction of wind shown by smoke drift | -21 |
| Light | 4 - 7 mph | Wind felt on face, leaves rustle | -31 |
| Gentle | 8 - 12 mph | Leaves/small twigs in constant motion. wind extends light flag | -35 |
| Moderate | 13 - 18 mph | Raises dust and loose paper; small branches are moved | -38 |

Outdoor Ventilation Adjustment: A: B:

A: B: Indoor Activities: On Table 1 on left, select the facility type and ACH (Air Changes per Hour) that best matches the activity location. If air is recirculated and filtered, add filter adjustments to the ACH. Next, find the adjustment that matches ACH and duration from table below. The durations used in Step 7 and Step 8 should match.

Table 1: Select ACH

| Medical | ACH |
|-----------------------------|------|
| General | 6 |
| Laboratory | 1.5 |
| Treatment room | 6 |
| Examination room | 6 |
| Retail | |
| Sales (except as below) | 1.5 |
| Barbershop | 1.5 |
| Hair and nail salons | 3.75 |
| Supermarket | 1 |
| Fast Food | 6 |
| Bars | 2-6 |
| Restaurants | 2-4 |
| Education | |
| Classrooms (ages 5 to 8) | 2 |
| Classrooms (age 9 plus) | 2 |
| Daycare (through age 4) | 2.5 |
| Multilevel assembly | 5 |
| Lecture hall (fixed seats) | 7 |
| Lecture classroom | 3 |
| Libraries | 1.5 |
| Music/theater/dance | 2.5 |
| Office | |
| Office space | 0.5 |
| Reception Area | 1.25 |
| Meeting/ Conference Rooms | 2 |
| Manufacturing | |
| Manufacturing Floor | 1.5 |
| Residential | |
| Homes with closed windows | 0.5 |
| Rooms with one open window | 1.75 |
| Homes with all open windows | 3 |
| Travel | |
| Aircraft | 20 |
| Trains/ Buses | 6 |
| Cars (Windows Closed) | 6 |
| Cars (Windows Open) | 10 |

Indoor Adjustment: A: B:

Find the ACH selected in Table 1
Time since activity or event started

| ACH | 5 min | 10 min | 15 min | 20 min | 45 min | 1 hr | 2 hr | 4 hr | 8 hr | 12 hr |
|------|-------|--------|--------|--------|--------|------|------|------|------|-------|
| 0.25 | -5 | -0 | +2 | +4 | +9 | +11 | +14 | +16 | +17 | +17 |
| 0.5 | -5 | -0 | +2 | +4 | +9 | +11 | +14 | +15 | +16 | +16 |
| 0.75 | -5 | -1 | +2 | +4 | +9 | +10 | +13 | +14 | +14 | +14 |
| 1 | -5 | -1 | +2 | +4 | +8 | +10 | +12 | +13 | +13 | +13 |
| 1.25 | -5 | -1 | +2 | +4 | +8 | +9 | +12 | +12 | +12 | +12 |
| 1.5 | -5 | -1 | +2 | +4 | +8 | +9 | +11 | +12 | +12 | +12 |
| 1.75 | -5 | -1 | +2 | +3 | +8 | +9 | +10 | +11 | +11 | +11 |
| 2 | -5 | -1 | +2 | +3 | +7 | +8 | +10 | +10 | +10 | +10 |
| 2.25 | -6 | -1 | +2 | +3 | +7 | +8 | +9 | +10 | +10 | +10 |
| 2.5 | -6 | -1 | +1 | +3 | +7 | +8 | +9 | +9 | +9 | +9 |
| 3 | -6 | -1 | +1 | +3 | +6 | +7 | +8 | +8 | +8 | +8 |
| 3.75 | -6 | -1 | +1 | +2 | +5 | +6 | +7 | +7 | +7 | +7 |
| 4 | -6 | -1 | +1 | +2 | +5 | +6 | +6 | +6 | +6 | +6 |
| 4.75 | -6 | -2 | 0 | +2 | +5 | +5 | +5 | +5 | +5 | +5 |
| 5 | -6 | -2 | 0 | +2 | +4 | +5 | +5 | +5 | +5 | +5 |
| 6 | -6 | -2 | 0 | +1 | +3 | +4 | +4 | +4 | +4 | +4 |
| 6.75 | -6 | -2 | 0 | +1 | +3 | +3 | +3 | +3 | +3 | +3 |
| 7 | -6 | -2 | 0 | +1 | +3 | +3 | +3 | +3 | +3 | +3 |
| 8 | -6 | -3 | -1 | 0 | +2 | +2 | +2 | +2 | +2 | +2 |
| 9 | -7 | -3 | -1 | 0 | +1 | +1 | +1 | +1 | +1 | +1 |
| 10 | -7 | -3 | -1 | -1 | +1 | +1 | +1 | +1 | +1 | +1 |
| 11 | -7 | -3 | -2 | -1 | 0 | 0 | 0 | 0 | 0 | 0 |
| 12 | -7 | -4 | -2 | -1 | -1 | -1 | -1 | -1 | -1 | -1 |
| 20 | -8 | -5 | -5 | -4 | -4 | -4 | -4 | -4 | -4 | -4 |
| 30 | -9 | -7 | -7 | -7 | -7 | -7 | -7 | -7 | -7 | -7 |

If air is recirculated and filtered, adjust ACH value upward

| Filter Type | Adjustment |
|---|---------------|
| If MERV 8 filters are used (Typical Home Filters) | Add +3 to ACH |
| If MERV 13 (Commercial) | Add +5 to ACH |
| If HEPA | Add +6 to ACH |

Step 9 If indoors, enter the room dimensions and the height of the ceiling.

Room Length (ft): A: B:

Room Width (ft):

Ceiling Height (ft):

Room Area (ft²): A: B:

For updates visit: <https://www.cov-irt.org/exposure-assessment-tool/>

For more information or questions go to: <http://www.signaturescience.com/contact>

Step 10 Calculate Adjustment to Local Community's Current Conditions

Average Daily Cases per 100,000 in the Last Week¹ x 5 x Average Days Infectious x Undiagnosed Factor for Area (Set to 5 if not known) = Active Infections per 100,000²

¹Lookup latest data here:

- <https://coronavirus.jhu.edu/us-map>
- https://covid.cdc.gov/covid-data-tracker/#cases_casesinlast7days
- <https://www.nytimes.com/interactive/2020/us/coronavirus-us-cases.html#map>

² Adjustment for community Infection rate is not made when exposure is due to exposure with people diagnosed with active COVID-19. N/A is shown.

Results Relative Exposure Ratio and near- and far-field contributions.

Ratios of Exposure to the Baseline Exposure Scenario³

A: B:
Near Field: Far Field: Near Field: Far Field:

| | Lower Exposure | Medium | High | Very High |
|---|----------------------|----------------------|----------------------|----------------------|
| A | <input type="text"/> | <input type="text"/> | <input type="text"/> | <input type="text"/> |
| B | <input type="text"/> | <input type="text"/> | <input type="text"/> | <input type="text"/> |

³The baseline exposure scenario is defined on Table 2 on next page. Results are equivalent to the baseline exposure when ratios of exposure are equal to 1. Results of 1 or greater are considered high exposures and higher risks can be inferred.

END Take actions to reduce exposure!

MEDIUM: When possible make adjustments to reduce the activity-related exposure. Consider mitigations such as wearing a mask increasing ventilation, increasing distancing, reducing people, or duration of the activity.
HIGH OR VERY HIGH: Make changes to reduce exposure. Rethink the event or activity. Combine multiple mitigations such as using N95 masks, increasing ventilation, holding event outside, greatly increasing distancing, reducing people, or shortening the duration.

Disclaimer Statement

By using this tool you agree to be bound by this disclaimer. This tool does not provide a prediction of the likelihood of infection or transmission of disease, nor does the tool provide advice or guarantee outcome. It solely provides a means for the user to estimate theoretical relative exposure. It is intended for informational purposes only and users should not rely on output for decision-making. Information is gathered from various sources and Signature Science, LLC, COV-IRT or its members are not responsible for errors or omissions in data. No warranties are given in relation to this tool and it is not guaranteed as fit for purpose.

A Simple-to-Use Tool (BETA Version) for Assessing Group Exposure to SARS-COV-2 based upon Mechanistic Factors, Group Behavior, and Infection Prevalence in the Community

The COVID-19 Exposure Assessment (CEAT) tool allows users to estimate a group-wide relative exposure to SARS-COV-2 associated with various activities. The tool facilitates understanding of the relative importance of factors that contribute to increased exposure. It incorporates both scientific principals and recent findings regarding virus characteristics to assess relative exposure based on the key mechanistic and epidemiological factors including: 1) the emission rate of virus, 2) viral aerosol degradation and removal, 3) duration of activity/exposure, 4) inhalation rates, 5) ventilation rates (indoors and outdoors), 6) volume of indoor space, 7) HVAC filtration removal efficiency, 8) mask use and effectiveness, 9) distance between people, 10) group population's adherence to public health guidance, 11) size of the group, and 12) prevalence of infection in the population. The model can be broadly applied to many situations (i.e., worker safety, public events, and schools). The model's results are aligned with the United States Occupational Safety and Health Administration (US OSHA) classifications of exposure risks [1], by benchmarking the exposure calculations to a baseline scenario that is considered high risk by US OSHA. We define the baseline scenario to represent a person (i.e., medical worker) who is exposed to a COVID-19 infected person. We apply assumptions to this scenario, addressing each of the factors in Table 2 to arrive at a baseline inhalation exposure dose value (exposure). The inhalation exposure for other scenarios are compared to the baseline by a ratio of the scenario exposure to the baseline exposure.

The starting point for the mechanistic inhalation exposure model is to use the relationship that defines inhalation exposure as:

$$\bar{E}_{mass} = \bar{C}_{AVE} \times Q_{inhalation} \times \Delta t \times P_{Exposed}$$

where $Q_{inhalation}$ is the inhalation rate, \bar{C} , is the average concentration, Δt is the duration of exposure, and $P_{Exposed}$ is the number of people exposed. \bar{E} is the total quantity of contaminant that enters the respiratory tract through the nose or mouth during inhalation over the duration of a potential exposure [2].

One critical variable that must be estimated by the model is the concentration of virus-containing aerosols that occurs as a result of the exhalation (i.e., breathing and singing) from people in close proximity. The underlying concentration model used in the model assesses both the contributions of concentration of virus due to the "nearness" of people (i.e., people in the "near field" whether indoors or outdoors) and the build up of concentration in a room over time (i.e., "far field") [3] [4]. Outdoors only the near field concentration contributions are used, since the far field is considered to be negligible [3]. Indoors, the far field result is approximately equivalent to the result that a well-mixed box assumption would provide.

For updates visit:

<https://www.cov-irt.org/exposure-assessment-tool/>

For more information or questions go to:

<http://www.signaturescience.com/contact>

Table 2. Exposure Factor Assumptions used in CEAT


| Exposure Factors | Range of Values used in the Model | Baseline Scenario Assumption |
|--|--|--|
| Emission rate of Infectious Aerosols Released through Breath | Range of viral RNA emissions based upon work by Buonanno, et al. [5][6] | 11.4 quanta per hour, associated with speaking while standing. |
| Inhalation Rate | Typical inhalation rates for adults at various activity intensities [7] | 0.46 cfm (13 liters per min), based on assumption of light intensity |
| Duration of Exposure | Varies between 5 minutes and 12 hours | 15 minutes, corresponding to the time defined by CDC contact tracing as a significant exposure [8] |
| Volume of Room or Space | Area varies between 100 ft ² (~10 m ²) and 5000 ft ² (~500 m ²). Height is assumed to be 9 ft (2.7 m). | 100 ft ² (~10 m ²), assuming a small room (e.g., a medical examination room) |
| Ventilation rates (Air changes per hour [ACH]) | Values based on published sources [9][10][11]. ACH impacted by outside air, recirculation rate, and effective ACH contribution from viral aerosol degradation and removal by deposition on surfaces. | 6 ACH, based on recommendation for medical treatment rooms [9]. Viral aerosol degradation/removal is assumed to be approximately 0.87 ACH for indoors (0.63 for viral degradation due to temperature and humidity and 0.24 due to surface deposition[12]). |
| Mask effectiveness | Range of mask effectiveness values based on published data for cloth, surgical, and N-95 masks [13][14] | Assume no masks are worn in the baseline scenario. |
| Distance between people | 15 ft (4.5 m), 10 ft (~3 m), 6 ft (~2 m), 3 ft (~1 m), and 1.5 ft (~0.5 m). | 3 ft (~1 m), based on estimated average distance during the event. |
| Number of people | Ranges from 2 to 250 people. | Two |
| Likelihood of Infectious persons present | Ranges over 6 orders of magnitude from the lowest (0.0001%) assumed for people adhering to strictly to public health guidance, to the highest (100%) for those known to be infected. | 100 percent likelihood of Infectious persons present (1 infected person is assumed) |

These concentration contributions are calculated for a person assumed to be at the center of a triangular grid where people are spaced equidistantly (based upon the distancing specified). We assume the concentration at the center is representative for all people in the group since: (1) each person's location is likely not static during the activity and (2) exposure is driven chiefly by the close-in sources (i.e., other people) and all people have close-in sources. The exposure calculation also incorporates the likelihood that people in the group are infected based on the prevalence of infection in the local community and assumed adherence to public health guidance, as well as the air changes per hour associated with the venue, the assumed activity level of individuals (affecting inhalation and exhalation rates) and the duration of the event.


The 11-step tool enables exposure calculations to be done in two ways (each are separately downloaded):

- Nomogram Version:** Uses a graphical nomogram with paper, pen and a straightedge.
- Adobe Reader® or Adobe Acrobat® Version:** Enabled by JavaScript®, this version uses dropdown menus and displays results in an interactive bar graph. **Note: Adobe Reader® or Adobe Acrobat must be used for the JavaScript to execute**

To accomplish the calculations, the tool uses a log-linear transformation, which allows the calculations of quotients and products to be made by adding and subtracting whole numbers. An order of magnitude decrease (increase) in exposure is achieved by a change of -17 (+17) in the adjustment factors.



COVID-19 International Research Team



Contributors:
 Brian Schimmoller, Signature Science, LLC
 Molly Isbell, Signature Science, LLC
 Som Dutta, Utah State University
 Afshin Beheshti, NASA Ames Research Center
 Tenley Archer, Algen Biotechnologies

The final results are relative exposure values, which are then categorized exposure into four bins, ranging from "Lower Exposure" through "Very High Exposure." Users can use the tool to assess the impact of changing scenario parameters (e.g., duration, number of people, spacing, mask use) to compare scenarios with the goal of lowering exposure and thereby lowering risk. Also, the results section shows the near field contribution to the total exposure from nearby people, as well as the resulting far field portion of the exposure from the build up of concentration in the room when indoors. The final result is the sum of the near field and far field exposure contributions.

List of References:

- OSHA. "Guidance on Preparing Workplaces for COVID-19", United States Occupational Safety and Health Administration (US OSHA), March 2020.
- US EPA. "Guidelines for Human Exposure Assessment." (EPA/100/B-19/001). Risk Assessment Forum, US EPA, October 2019
- "Toxic Risk Symposium: Estimating Exposure for On-Site Worker Health Risk Estimates" October 2014. <https://www.aqmd.gov/home/rules-compliance/compliance/vocs/exempts/toxic-symp>.
- Jaycock, M et al., "The Daubert Standard as Applied to Exposure Assessment Modeling Using the Two-Zone (NF/FF) Model Estimation of Indoor Air Breathing Zone Concentration as an Example." Journal of Occupational and Environmental Hygiene 8, no. 11 (November 2011): D114-122. <https://doi.org/10.1080/15459624.2011.624387>.
- Buonanno, G., et al., "Quantitative Assessment of the Risk of Airborne Transmission of SARS-CoV-2 Infection." Environment International 145 (December 2020): 106112.
- Buonanno, G., et al., "Estimation of Airborne Viral Emission: Quanta Emission Rate of SARS-CoV-2 for Infection Risk Assessment." *Environment International* 141 (August 2020)
- US EPA, ORD. "Exposure Factors Handbook Chapter 6." Reports and Assessments. US EPA, September 18, 2015. <https://www.epa.gov/expobox/exposure-factors-handbook-chapter-6>.
- CDC. "Public Health Guidance for Community-Related Exposure." CDC. <https://www.cdc.gov/coronavirus/2019-ncov/php/public-health-recommendations.html>.
- CDC. "Appendix B. Air Guidelines for Environmental Infection Control in Health-Care Facilities," July 22, 2019. <https://www.cdc.gov/infectioncontrol/guidelines/environmental/appendix/air.html>.
- ANSI/ASHRAE Standard 62.1-2019, "Ventilation for Acceptable Indoor Air Quality," 2020. <https://www.ashrae.org/technical-resources/bookstore/standards-62-1-62-2>.
- Howard-Reed, Cynthia, et al., "The Effect of Opening Windows on Air Change Rates in Two Homes." Journal of the Air & Waste Management Association 52, no. 2 (February 1, 2002): 147-59. <https://doi.org/10.1080/10473289.2002.10470775>.
- CIRES. "COVID-19 Airborne Transmission Tool Available." Text. CIRES, June 25, 2020. <https://cires.colorado.edu/news/covid-19-airborne-transmission-tool-available>.
- Mueller, Amy V., et al., "Quantitative Method for Comparative Assessment of Particle Removal Efficiency of Fabric Masks as Alternatives to Standard Surgical Masks for PPE." *Matter* 3, no. 3 (September 2020): 950-62
- CDC. "Scientific Brief: Community Use of Cloth Masks to Control the Spread of SARS-CoV-2, Centers for Disease Control and Prevention, November 2020. <https://www.cdc.gov/coronavirus/2019-ncov/more/masking-science-sars-cov-2.html>.

Disclaimer Statement:

By using this tool you agree to be bound by this disclaimer. This tool does not provide a prediction of the likelihood of infection or transmission of disease, nor does the tool provide advice or guarantee outcome. It solely provides a means for the user to estimate theoretical relative exposure. It is intended for informational purposes only and users should not rely on output for decision-making. Information is gathered from various sources and Signature Science, LLC, COV-IRT or its members are not responsible for errors or omissions in data. No warranties are given in relation to this tool and it is not guaranteed as fit for purpose.